

POLAND/Physical Chemistry - Surface Phenomena, Adsorption,
Chromatography, Ion Exchange.

B.

Abs Jour : Ref Zhur - Khimiya, No 14, 1958, 46139

adsorption of the air itself and considerably less in
the A of CS₂. The presence of air does not influence
the A degree of CS₂, because the apparent change in the
adsorption properties of carbon are caused by the de-
sorption of air.

Card 2/2

POLAND / Chemical Technology, Chemical Products and
Their Application, Part 4. - Artificial and
Synthetic Fibers. H

Abs Jour: Ref Zhur-Khimiya, No 18, 1958, 63031.

Author : ~~Mieczyslaw Wronski~~

Inst : Lodz University.

Title : Complete Analysis of Viscose.

Orig Pub: Zesz. nauk, Univ. odzk., 1957, Ser. 2, No 3,
159 - 165.

Abstract: Cellulose xanthogenate (I) is deposited from
viscose with saturated NaCl solution. Na_2CS_3
and Na_2CS_4 are determined photometrically in
the filtrate. Na_2S and $\text{Na}_2\text{S}_2\text{O}_3$ are determined
by direct iodometric titration, NaOH and Na_2CO_3
are determined acidimetrically. The con-
centration of I is found from the difference

Card 1/X2

POLAND / Chemical Technology, Chemical Products and
Their Application, Part 4. - Artificial and
Synthetic Fibers.

H

Abs Jour: Ref Zhur-Khimiya, No 18, 1958, 63031.

Abstract: between the iodine amounts consumed by the
titration of the initial viscose and of the
filtrate. The complete analysis takes 20
min. or less.

Card 2/2

43

POLAND/Physical Chemistry. Surface Phenomena, Adsorption.
Chromatography, Ion Exchange.

B

Abs Jour: Ref Zhur-Khim., No 15, 1958, 49752.

Author : Wronski, Mieczyslaw.

Inst : Lodz University.

Title : Sorption of Carbon Disulfide by Alkali Cellulose.

Orig Pub: Zesz. nauk. Uniw. lodzk., 1957, Ser. 2, No 3, 167-170.

Abstract: Study of temperature dependence of the adsorption of
CS₂ vapor by alkali cellulose. CS₂ adsorption curves
show a minimum at a temperature of about 14° which
indicates chemical and physical adsorption. --
Author's summary.

Card : 1/1

WRONSKI, MIECZYSLAW

H.

POLAND/Artificial and Synthetic Fibers.

Abs Jour : Ref Zhur - Khimiya, No 19, 1958, 66205

Author : Wronski Mieczyslaw

Inst :

Title : An Investigation of the Penetration of a Precipitating Bath Through Layers of Viscose.

Orig Pub : Zesz. nauk. Univ. lodzk., 1957, Ser. 2, No 3, 171-175.

Abstract : By means of a glass electrode, the rate of penetration of a precipitating bath through layers of viscose was investigated. The derived pH-time experimental graph of the contact of a viscose layer, found on the electrode, with the precipitating bath, possess three small curves. The first corresponds to the neutralization of NaOH and the formation of Na_2SO_4 and Na_2CO_3 ; the second, to the decomposition of these salts; the third, to the decrease of the concentration of hydrogen ions. Proceeding from the assumption that through the layer formed of

Card 1/2

POLAND/Physical Chemistry. Kinetics. Combustion. Explosions.
Topochemistry. Catalysis.

B

Abs Jour: Ref Zhur-Khim., No 15, 1958, 49623.

Author : Wronski, Mieczyslaw.

Inst : Lodz University.

Title : Kinetics of Decomposition of Sodium Ethyl Xanthogenate
in Caustic Alkali.

Orig Pub: Zesz. nauk. Uniw. lodzk., 1957, Ser. 2, No 3, 177-185.

Abstract: Determination of the correlation between rate of
decomposition of $C_2H_5OCSSNa$, with formation of
 Na_2S and Na_2CS , and concentration of caustic
alkali, temperature, and the presence of Na_2S .
Decomposition of xanthogenate occurs according to
two distinct schemes: $ROSS^- = RO^- + CS_2$ and $ROCSS^- +$

Card : 1/2

POLAND/Physical Chemistry. Kinetics. Combustion. Explosions.
Topochemistry. Catalysis.

D

Abs Jour: Ref Zhur-Khin., No 15, 1958, 49623.

$\text{OH}^- = \text{ROH} + \text{CS}_2\text{O}^{2-}$. Rate of decomposition of xantho-
genate is defined by the equation: $-\text{dx}/\text{dt} = k_1x +$
 $k_2x (\text{NaOH})^2$. -- Author's summary.

Card : 2/2

26

POLAND / Analytical Chemistry. Analysis of Organic Substances.

E-3

Abs Jour: Ref Zhur-Khimiya, No 1, 1959, 1029.


Author : Wronski, M. Univ. Lodz, Poland

Inst : ~~Not given.~~

Title : The Titration of Sulfides With o-Hydroxy Mercurobenzoic Acid.

Orig Pub: Chem analit., 1957, 2, No 4, 385-386.

Abstract: The titrimetric method for the determination of S^{2-} , in the presence of SO_3^{2-} , $S_2O_3^{2-}$ and xanthogenates is described, the method being based on the reaction:



From 0.1 to 0.5 millimoles of Na_2S is dissolved

Card 1/3

POLAND / Analytical Chemistry. Analysis of Organic Substances.

E-3

Abs Jour: Ref Zhur-Khimiya, No 1, 1959, 1029.

Abstract: in 150 millimeters of water from which oxygen has been removed previously by the addition of Na_2SO_3 , then 5 milliliters of 0.5 N NaOH is added followed by a few drops of 0.1% sodium nitroprusside solution, and the mixture is titrated with 0.05 M solution of 6-hydroxy mercurobenzoate of Na (I) until the disappearance of the violet color. The solution of I is prepared by dissolving o-hydroxy mercurobenzoic anhydride in a 0.25 N NaOH solution. A titre of the solution obtained is determined iodometrically; for that purpose 10 millimeters of concentrated sulfuric acid and 10 millimeters of 0.1N iodine solution are added to 10 millilit-

Card 2/3

28

The trithiocarbonate formation in the xanthate reaction
Mieczyslaw W. Wronski
The effect of the temperature on the
rate of the reaction was studied. The results are
discussed in terms of the mechanism of the reaction.

Wronski

POLAND / Laboratory Equipment, Apparatus; Their Theory,
Construction and Application.

F

Abstr Jour : Ref Zhur - Khim., No 10, 1958, No 32280

Author : Jozef Chrzaszowski, Mieczyslaw Wronski.

Inst : -

Title : Simple Determination Method of Isotherm of Vapor Adsorption
on Solid Substances.

Orig Pub : Roczn. chem., 1957, 31, No 1, 297-299

Abstract : A simple apparatus for measuring isotherms of vapor
adsorption is described. The apparatus consists of a gas
burette connected with a Hg manomotor, vacuum installation
and two vessels with faucets for the adsorbent and adsorbed
substance. Computation equations are presented.

Card 1/1

WROŃSKI, M.
 POLAND / Laboratory Equipment, Apparatus, Their Theory, F
 Construction and Application.

Abs Jour: Ref Zhur-Khimiya, No 18, 1958, 60749.

Author : Mieczyslaw Wronski.

Inst :

Title : Effect of Width of Spectral Zone on Photometric Measurements.

Orig Pub: Roczn. chem., 1957, 31, No 1, 309-313.

Abstract: The effect of the polychromaticity of light on the extinction (E) measurements was computed. Making some simplifying assumptions, the author receives for the value of E: $E = \log[2.3(\sum_2 - \sum_1)k/(10^{-k\sum_1} - 10^{-k\sum_2})]$, where k is the product of the solution concentration and the thickness of the absorbing layer, and \sum_2 and \sum_1 are the E factors at the

Card 1/2

1

✓ Titration of hydrogen sulfide and other sulfides with organic mercury compounds. (Mieczysław Wroński and Philipp Burkart. *Faserforsch. u. Textiltech.* 9, 38-7 (1958).—A 0.05N soln. of 2-HOOCCH₂HgOH is used for the titration of S²⁻ or HS⁻, a colorless 0.5% soln. of thiofluorescein (the dimercapto analog of fluorescein) in 0.1N NaOH being used as indicator. In the presence of S²⁻, a sharp change to dark blue indicates the end of the titration. The presence of large amts. of I⁻, NCS⁻, or Cl⁻ does not disturb the reaction. Since CN⁻, xanthates, or S₂O₃²⁻ disturb the reaction, the use of dithione as indicator is recommended when xanthates or S₂O₃²⁻ is present in larger amts. Polysulfides are also titrated, but the addn. of SO₃²⁻ is indicated for better end-point detn. The method shows excellent correlation with the complexometric detn. and is also suitable for the H₂S analysis of air or exhaust gases. Paul D. Burgauer

WRONSKI, M.

Organic mercury compounds in chemical analysis. Młeczysław Wronski (Univ. Łódź, Poland). *Zeszyty Nauk. Univ. Łódzkiego*, Ser. II No. 4, 181-93 (1958) (English summary).

Compds. of the type $RHgOH$, sol. in alkalies owing to the presence of OH or COOH groups (mainly $o-HOC_6H_4HgOH$ (I)), were used in the volumetric detn. of some S compds. in the presence of color indicators. The titrant (0.05M) was prepd. by dissolving I in 0.2N NaOH and standardized with $Na_2S_2O_3$ or Na_2S . To det. S^{--} , 100 ml. of a soln. contg. $5 \times 10^{-3} - 2 \times 10^{-2}$ g. H_2S was treated with 5 ml. N KOH or NaOH, 1 ml. of 0.1% dithizone in EtOH, and titrated with I until the yellow color turned red; SO_3^{--} , $S_2O_3^{--}$, CNS^- , thiourea, and moderate quantities of xanthates (II) (unlike mercaptans, thiocarbonates (III), and CN^-) did not interfere with the detn. Dithiocarbamates were titrated in the presence of dithiofluorescein (IV) (0.05% soln. in 0.01N NaOH with I-Na added to full decolorization) until the blue color completely disappeared. Na_2S_2 can be titrated like Na_2S , but more reliably after reln. to Na_2S by heating with 10% Na_2SO_3 and then titrating as usual (result A); to det. Na_2S and Na_2S_2 in mixts. a 2nd titration was necessary. The soln. treated with excess I, heated to boiling, cooled, treated with excess Na_2S , and titrated as usual (result B); Na_2S content = $2.5(A - 0.0B)$ and Na_2S_2 content = $1.5(B - A)$. III were detd. most conveniently by titrating with Na_2S the excess of

I in the presence of IV, or by direct titration with $Hg(NO_3)_2$ in slightly alk. medium; S^{--} interfered with this detn., but the sum of S^{--} and III was detd. by adding a known quantity of I prior to titration. To det. $S_2O_3^{--}$, a sample equiv. to 5-10 ml. of 0.01N $Na_2S_2O_3$ was treated with 20 ml. acetate buffer, dild. to 100 ml. treated with 1 ml. 0.1% EtOH soln. of diphenylcarbazone (V), and titrated to

a blue color; SO_3^{--} , I^- , CNS^- , and thiourea obstructed the detn. Thiophenols, II, and mercaptobenzthiazoles were titrated with bis(hydroxymercuri)thymol (VI) or $Hg(NO_3)_2$ in slightly alk. medium, and in the presence of V; the detn. was obstructed by I^- , $S_2O_3^{--}$, SO_3^{--} , and thiourea, but not by SO_3^{--} , Cl^- , and small quantities of CNS^- , Br^- , or NH_4^+ . To det. II in the presence of S^{--} and III, the soln. was acidified with N HCl and then made alk. with N NaOH; transforming thereby III into S^{--} , the sum of which was detd. by I titration; in a sep. sample all 3 components were detd. iodometrically, and II was calcd. as the difference. The system $Na_2S-Na_2S_2O_3-Na_2SO_3$ was titrated with VI, S^{--} in the presence of IV, and then, after adding NH_4Cl , thiosulfates in the presence of V; SO_3^{--} was detd. by addnl. iodometric titration. Hg derivs. of phenolphthalein and fluorescein proved sensitive to some S compds.; this property may be useful in colorimetric assays. Titration of S compds. with I can be reversed and used for detg. Hg in org. compds.; IV was preferred in such detns. and used in excess which should be back-titrated with standard $Hg(NO_3)_2$. All indicators are described in detail; monothiodifluorescein, dimercaptophenolphthalein, and di-2-naphthylthiocarbazon were used besides those mentioned above. Dissocn. consts. of the following adducts were detd. photometrically: $I + IV$, 2.30×10^{-4} in 0.04N, and 6.25×10^{-4} in 0.002N NaOH; $1.5I + V$, 2.08×10^{-4} in 0.08N, and 1.58×10^{-4} in 0.002N NaOH; $I + V$ - 6.1×10^{-4} at pH 5.0. J. Lange

B

Country : Poland
 Category : Physical Chemistry - Kinetics. Combustion. Ex-
 plosions. Topochemistry. Catalysis. 45131
 Abs. Jour : RZhKhim., No 13, 1959
 Author : Wronski, M.
 Institut. : Not given
 Title : The Kinetics of the Reaction of Sodium Hydrate
 with Carbon Disulfide
 Orig. Pub. : Roczniki Chem, 32, No 4, 849-861 (1958)
 Abstract : The author has investigated the kinetics of the
 reaction of NaOH with CS₂ as a function of the
 concentration of the alkali solution (0.45-5.00 M)
 at 15 and 25°. The effect of the addition of
 Na₂SO₃ (0.125 M Na₂SO₃ in 1 M NaOH) on the reac-
 tion rate was studied. The concentration of the
 reaction products, Na₂S and Na₂CS₃, was determined
 by amperometric titration. It has been found that
 the rate of NaOH consumption is described by the
 equation:

$$-d[\text{NaOH}]/dt = k_1 [\text{NaOH}][\text{CS}_2]$$

 and that the rate of Na₂CS₃ formation follows the
 equation:

Card: 1/3

Country : Poland

B

Category :

Abs. Jour :

45151



Author :

Institut. :

Title :

Orig Pub. :

Abstract : $d[\text{Na}_2\text{CS}_3] = k_2[\text{Na}_2\text{S}][\text{CS}_2] + k_3[\text{NaOH}][\text{CS}_2]$ (1)
The value of k_2 increases with increasing initial concentration of the NaOH solution, as can be expected on the basis of current theories on the solvation of the S^{2-} ion. The addition of Na_2SO_3 has no effect on the reaction rate. Notwithstanding the earlier proposed reaction scheme:



(BZhKhim, No 2, 1953, 1574), the formation of Na_2CS_3 according to equation (1) proceeds in a more complicated way. In the opinion of the author the initial step in the reaction of NaOH with

Card: 2/3

Country : Poland
Category :

B.

Abs. Jour : 45131



Author :
Institut. :
Title :

Orig Pub. :

Abstract : CS_2 involves the formation of the ion CS_2OH^- by the reaction



The latter ion dissociates in two ways:



and



For reaction (2) values of $\Delta H^\ddagger = 21.4 \text{ kcal/mol}$ and $\Delta S^\ddagger = 0 \text{ e.u.}$ have been obtained.

C. Polotnyuk

Card: 3/3

P/012/59/004/03/05/020

82240

5.3200

AUTHOR:

Wroński, M.

TITLE:

The Kinetics of the Xanthate Reaction of Starch, Cellulose and Sodium Alginate

PERIODICAL:

Societas Scientiarum Lodziensis Acta Chimica, 1959, Vol 4, pp 47 - 54

TEXT:

Owing to the great technological importance of the process of cellulose sulphidizing, numerous investigations of this reaction have been carried out before, but methods used were such that clear interpretation of results was impossible. This was so, because the reactions between alkali-cellulose and gaseous carbon disulphide are rather complicated owing to the adsorption and diffusion which obscures the real kinetic course. In this investigation, measurement of the sulphidizing rates in a single phase arrangement were carried out with a constant concentration of carbon disulphide. Because of this, the interpretation of results was easy. The process of formation of cellulose and starch xanthates, sodium salt of alginic acid and of some by-products was examined; the speed of cellulose xanthate decomposition in NaOH solutions was also investigated. From the results

Card 1/2

OK

82240

P/012/59/004/03/05/020

The Kinetics of the Xanthate Reaction of Starch, Cellulose and Sodium Alginate

obtained the speed of reaction constants was calculated. In the case of starch it was found that introduction of the second xanthate group into the glucose ring is much more difficult than in the case of cellulose. Sodium hydroxide and sodium chloride both suppress the speed of cellulose xanthate decomposition. There are 4 figures, 2 tables and 6 references: 3 Polish, 2 German and 1 English.

ASSOCIATION: Katedra Technologii Chemicznej Uniwersytetu Łódzkiego (Lodz University, Department of Chemical Technology)

PRESENTED: March 11, 1960

Card 2/2

P/012/59/004/03/06/020

5.1200

AUTHOR: Wroński, M.

82241

TITLE: The Kinetics of the Xanthate¹ Reaction of Allyl and Furfuryl Alcohol, Glycolic Acid and Methylene Glycol¹

PERIODICAL: Societas Scientiarum Lodziensis Acta Chimica, 1959. Vol 4, pp 55 - 63

TEXT: The author presents the continuation of his investigations concerning the mechanism of xanthate reaction, generally expressed by the following equation: $\text{ROH} + \text{NaOH} + \text{CS}_2 = \text{ROCSSNa} + \text{H}_2\text{O}$. The process of formation and decomposition of xanthates of aliphatic mono- and polyalcohols, glucose and saccharose has been investigated by the author before. This report presents the results of investigations concerning the influence of some groupings on the course of xanthate reaction. Apparently no investigations of xanthation of such alcohols has been made yet so far. The process of xanthate and by-product formation during the sulphidizing of allyl- and alpha furfuryl alcohols, glycolic acid and methylene glycol at 15° and 25° as well as the speed of allyl xanthate decomposition at 55° and 65°C were investigated. From results obtained the speed of reaction constants

Card 1/2

P/012/59/004/03/06/020

The Kinetics of the Xanthate Reaction of Allyl and Furfuryl Alcohol, Glycolic Acid and Methylene Glycol

82241

was calculated. Allyl alcohol and glycolic acid form fairly stable xanthates, while methylene glycol xanthate hydrolyses instantaneously after formation. In conformity with this, methylene glycol catalyses the hydrolysis of carbon disulphide in the presence of NaOH. During the process of alpha-furfuryl alcohol sulphidizing the monothiocarbonate appears in quantities largely exceeding the amount of by-products formed. This could be explained if one admits that sulphur can extrude oxygen from the furan ring. No reaction has been observed between sodium phenolate and carbon disulphide. There are 5 figures, 1 table and 7 references: 1 English and 6 Polish.

ASSOCIATION: Katedra Technologii Chemicznej Uniwersytetu Łódzkiego (Łódź University, Department of Chemical Technology)

PRESENTED: March 11. 1959

Card 2/2

WRONSKI, M.

✓ Kinetics of xanthation of polybasic alcohols. Mieczysław
Wronski (Univ. Łódź, Poland). *Zeszyty Nauk. Chem.*
Łódzkiego, Ser. II, No. 3, 191-202 (1969).—Xanthation of
ethylene glycol (I), glycerol (II), D-glucose (III), and tri-
ethylene glycol (IV) was investigated kinetically as earlier
methylene glycol (IV) was investigated kinetically as earlier
(preceding abstr.). The following data were found in solns.
contg. 50 g./l. of the alc. in 0.5N NaOH (compd., temp.,
 k_1 , k_2 , k_3 , and k_4 given): I, 15°, 0.10, 0, 2.4×10^{-3} , 0,
 5×10^{-3} ; I, 25°, 1.1, 0, 6.6×10^{-3} , 0, 8×10^{-3} ; II,
15°, 0.79, 0, 0.16, 0, 0.8×10^{-3} ; II, 25°, 1.0, 0, 0.36, 0,
 2.4×10^{-3} ; III, 15°, 0.43, 2.5×10^{-3} , 0, 2.2×10^{-3} ,
 2.0×10^{-3} ; III, 25°, 1.3, 1.2×10^{-3} , 0, 0.14, 6.1×10^{-3} ;
IV, 15°, 0.384, —, —, —; IV, 25°, 0.915, —, —, —;
IV, 40°, —, 9.4×10^{-4} , —, 5×10^{-4} , —; IV, 50°, —,
 1.7×10^{-3} , —, 1.4×10^{-3} , —. Decompu. of the xan-
thates of I and II followed the equation $-dx/dt = k_{1x}$
[NaOH], whereas those of III and IV decompd. according to
 $-dx/dt = k_{2x} + k_{4x}$ [NaOH]². Several graphs were repro-
duced and detailed anal. procedure was given. J. Lange

3
17-jaj/148

✓ Synthesis of 2-(o-hydroxyphenyl)benzoxazole. Mieczysław Wroński (Univ. Łódź, Poland). *Roczniki Chem.* 33, 809-10 (1959) (English summary). — P_2O_5 (28 g.) is added during 2 hrs. to 15 g. salicylic acid and 10 g. o-aminophenol with intensive mixing and keeping the temp. at 160-80°. The product is heated with H_2O , filtered, the ppt. dissolved in 100 ml. hot EtOH, again pptd. by diln. with H_2O , and purified further (Walter and Freiser, *C.A.* 46, 9011f). The yield of 2-(o-hydroxyphenyl)benzoxazole, m. 122-4°, is 60%.
A. Kreglewski

3
4E 3 d
122 (NB)

Card 1/1

aht

99

WRONSKI, Mieczyslaw

Kinetics of xanthate reaction of simple alcohols. Roczniki chemii 33
no.4/5:1061-1069 '59. (KRII 9:9)

1. Zaklad Technologii Chemicznej Uniwersytetu, Lodz.
(Xanthates) (Carbon disulfide) (Methanol)
(Ethyl alcohol) (Propyl alcohol) (Butyl alcohol)
(Isopropyl alcohol)

WROŃSKI, M.

Distr: 4E2o(j)

The kinetics of decomposition of xanthogenates in sodium hydroxide solutions. Mieczysław Wronski (Univ. Łódź, Poland). *Roczniki Chem.* 33, 1071-80 (1959) (German summary).—The rate of decomn. of Me, Et, Pr, iso-Pr, and Bu xanthogenates at 65 and 75° is expressed by the following equation as a function of NaOH concn.: $-dx/dt = k_1x +$

$k_2[NaOH]^4$. The ratio k_1/k_2 is about 2 for the Me ester and about 3 for others. The values of the consts. decrease: Me > Et > Bu > iso-Pr. For NaOH concns. below 0.3N the reaction can be expressed by $-dx/dt = k_1x$.

A. Kreglewski

3
1-gg(NB)

gg

WRONSKI, M.

Kinetics of the xanthate reaction of simple alcohols.
M. Wronski (Univ. Lodz, Poland). Z. physik. Chem.
(Leipzig) 211, 113-17 (1959). The xanthate reaction of
simple alcs. is assumed to proceed according to: $\text{ROH} +$
 $\text{OH}^- \rightarrow \text{RO}^- + \text{H}_2\text{O}$; $\text{RO}^- + \text{CS}_2 \rightarrow \text{ROCSS}^-$. If dx/dt
 $= k_1[\text{RO}^-][\text{CS}_2]$, $K = [\text{RO}^-]/[\text{ROH}][\text{OH}^-]$, $\xi = [\text{S}^{--}] +$
 $[\text{CS}_2^{--}]$, and $a =$ initial concn. of ROH, then $\ln(a/(a-x))$
 $= k_1 K \xi / k_2$. The values obtained by aid of this equation
conform well with those found in the literature. P. E.

3
292 (18)

99

WRONSKI, M.

4230
11/54

Maximum reaction velocity of carbon disulfide in sodium hydroxide. M. Wronski (Univ. Lodz, Poland). Z. physik. Chem. (Leipzig) 211, 113-20 (1959); cf. C.A. 53, 18311b. The reaction velocity of CS₂ in NaOH shows a distinct maximum that is caused by the slope of the product (OH⁻·XCS₂) as a function of the NaOH ((CS₂) = satn. concn. of CS₂ in NaOH). This max. can be calcd. by aid of the formula: (OH⁻·XCS₂) = y = G(CS₂) = G × 10⁴ - A, wherein G = (NaOH) in mole/l., m = 0.163, and A = a temp.-dependent const. If for a given end concn. of NaOH, G₁, the initial concn., G₀, is so to be chosen that the reaction runs off in a min. of time, G₀ can be detd. by G₀ = (1/2.3m) + Friedrich, Epstein.

3

17c

WRONSKI, Mieczyslaw

Indirect mercurimetric determination. Chem anal 5 no.1:101-107 '60.
(EEAI 9:11)

1. Zaklad Technologii Chemicznej Uniwersytetu, Lodz.
(Mercurimetry)

WRONSKI, Mieczyslaw

Determination of small amounts of silver and mercury by using
thiofluorescein. Chem anal 5 no.2:289-291 '60. (EEAI 10:3)

1. Zaklad Technologii Chemicznej Uniwersytetu, Lodz.
(Silver) (Mercury) (Thiofluorescein)

WRONSKI, Mieczyslaw

Argentometric determination of cyanide with a thiofluorescein indicator.
Chem anal 5 no.2:293-296 '60. (EEAI 10:3)

1. Zaklad Technologii Chemicznej Uniwersytetu, Lodz.
(Argentometry) (Cyanides) (Thiofluorescein)

WRONSKI, Mieczyslaw

The indirect colorimetric determination of sulfide and cyanide
with the aid of thiofluorescein. Chem anal 5 no.3:457-460 '60.
(EEAI 10:8)

1. Zaklad Technologii Chemicznej Uniwersytetu, Lodz.
(Colorimetry) (Sulfides) (Cyanides) (Thiofluorescein)

WRONSKI, Mieczyslaw

Titration of mercury and nickel salts with cysteine solution.
Chem anal 5 no.3:511-512 '60. (KEAI 10:8)

1. Zaklad Technologii Chemicznej Uniwersytety, Lodz.
(Mercury) (Nickel) (Cysteine) (Solutions)

WRONSKI, Mieczyslaw

Volumetric determination of trace amounts of copper with oxin blue.
Chem anal 5 no.4:597-599 '60. (EEAI 10:9)

1. Department of Chemical Technology, Lodz.

(Copper) (Oxin blue)

WRONSKI, Mieczyslaw

Rapid determination of mercury compounds in crude phenylmercury acetate. Chem anal 5 no.4:601-604 '60. (EEAI 10:9)

1. Department of Chemical Technology, University, Lodz.

(Mercury) (Phenylmercury acetate)

WRONSKI, Mieczyslaw

Mercurimetric determination of styrene, acrylonitrile and methyl
acrylate. Chem anal 5 no.5:823-826 '60. (ZEAI 10:9)

1. Department of Chemical Technology, University, Lodz.

(Mercurimetry) (Styrene) (Acrylonitrile)
(Methacrylate)

WRONSKI, Mieczyslaw

The influence of acids on the rate of mercurization of phenol and aniline. Roczniki chemii 34 no.3/4:947-952 '60. (EEAI 10:3)

1. Katedra Technologii Chemicznej Uniwersytetu, Lodz.
(Acids) (Phenol) (Aniline) (Mercury)

WRONSKI, Mieczyslaw

Desulfurating titration of organic sulphur compounds. Chem anal 6
no.5:869-876 '61.

1. Department of Chemical Technology, University, Lodz.

WRONSKI, Mieczyslaw, Doc. Dr. inz. (Lodz, Nowotni 18)

Mercurimetric determination of sulfur compounds applying acrylnitrile
as selective masking agent. Acta chimica Hung 28 no.1/3:303-309
'61. (EEAI 10:9)

1. Institut fur Chemische Technologie der Universitat Lodz, Polen.

(Mercurimetry) (Sulfur) (Acrylonitrile)

WRONSKI, Mieczyslaw

Accuracy of titration of sulfide with the sodium salt of
o-hydroxymercuribenzoic acid. Nauki matematyczne Lodz
no.10:205-210 '61.

1. Department of Chemical Technology, University, Lodz.

WRONSKI, Mieczyslaw; HAZEK, Lucyna

Kinetics of the hydrolysis of Phenyl isothiocyanate in solutions of sodium hydroxide. Nauki matematyczne Lodz no.12:155-162 '62.

1. Katedra Technologii Chemicznej, Uniwersytet, Lodz.

WRONSKI, Mieczysław

Speedy determination of mercury in mercury preparations.
Chem anal 7 no.4:821-826 '62.

1. Department of Chemical Technology, University, Lodz.

WRONSKI, Mieczyslaw

Determination of thioglycolic acid in the presence of sulfide, sulfite and thiosulfate. Chem anal 7 no.4:851-854 '62.

1. Department of Chemical Technology, University, Lodz.

WRONSKI, Mieczyslaw

Microdetermination of sulfides and thiourea in thiocyanates.
Chem anal 7 no.5:1009-1010 '62.

1. Katedra Technologii Chemicznej, Uniwersytet, Lodz.

WRONSKI, Mieczyslaw

Determination of mercuric acid in phenylmercuric acetate. Chem
anal 7 no.5:1011-1012 '62.

1. Katedra Technologii Chemicznej, Uniwersytet, Lodz.

S/081/63/000/003/008/036
B144/B186

AUTHOR: Wronski, Mieczyslaw

TITLE: Desulfurating titration of organic sulfur compounds

PERIODICAL: Referativnyy zhurnal. Khimiya, no. 3, 1963, 141, abstract
3G159 (Chem. analit. (Polska), v. 6, no. 5, 1961, 869-876
[Eng.; summary in Pol.])

TEXT: When o-hydroxy mercury benzoic acid (OA) acts on compounds contain-
ing CS groups in alkaline medium, compounds of the type R-Hg-S-Hg-R are
formed. It is suggested that this reaction be used for determining
compounds containing hydrolyzable sulfur (e. g. thiourea, thioacetamide)
by direct titration of the solutions with OA. Making use of the differences
in reaction rates, it is possible by this method to determine the sulfur
compounds separately in the presence of others. The reaction rate
increases with increasing concentration of the base and rising temperature.
The compounds studied can be arranged in the following order according to
the decreasing rate of reaction with OA: phenyl monothiocarbamate (I),
ethyl dithiocarbamate (II), benzyl dithiocarbamate (III), phenyl dithio-
carbamate (IV), dithiocarbamate (V), β -amino ethyl dithiocarbamate (VI),
Card 1/4

Desulfurating titration of organic ...

S/081/63/000/003/008/036
B144/B186

β -hydroxy ethyl dithiocarbamate (VII), diphenyl thiourea (VIII), rubeanic acid (IX), ethyl monothiocarbamate (X), rhodanine (XI), β -naphthyl thiourea (XII), thiourea (XIII), thiosemicarbazide (XIV), thioacetamide (XV), cellulose xanthate (XVI), trithiocarbonate (XVII), methyl xanthate (XVIII), bis-hydroxy ethyl dithiocarbamate (XIX), mercapto thiazoline (XX), dithiocarbazinate (XXI), phenyl dithiocarbazinate (XXII), ethyl xanthate (XXIII), diethyl dithiocarbamate (XXIV), o-phenylene thiourea (XXV), mercapto benzothiazole (XXVI), ethylene thiourea (XXVII), mercapto thioketo thiodiazole (XXVIII), thiosulfate (XXIX), thiocyanate (XXX). For determining I, and V - X, 5 ml 1 N NaOH solution, water or (in the case of insoluble compounds) CH_3OH up to a volume of 30 to 50 ml are added to the sample, and the mixture is titrated with 0.001 - 0.05 N OA solution, as described previously (RZhKhim, 1960, no. 20, 80867). As indicator is added 0.5 ml of 20 mg thiofluorescein (XXXI) dissolved in several ml of 1 N NH_4OH solution, diluted to a volume of 50 ml by 0.05 N solution of ethylene diamine tetraacetic acid, or 0.2 ml 0.1% solution of dithizone (XXXII) in $\text{C}_2\text{H}_5\text{OH}$. In the first case titration is carried out till the blue color disappears; in the second case till the yellow color

Card 2/4

Desulfurating titration of organic ...

S/081/63/000/003/008/036
B144/B186

turns purple. Titration is carried out at 30 - 40°C. Samples II-IV are prepared in the same manner; 5 - 20 ml toluene is added to the solution and titrated at 20°C in the presence of XXXI, as long as the blue color does not disappear for at least 30 sec. Samples XI - XV are dissolved in 5 ml 1 N NaOH solution, diluted to a volume of 25 ml and titrated with 0.05 N OA solution at 80 - 90°C in the presence of XXXI. In the titration of XII - XV 1 - 2 ml excess OA solution is added; after some minutes 25 ml cold water and 2 ml 0.1 N Na₂S solution containing 2% Na₂S and 1% NaOH, are added, and the Na₂S excess is titrated with OA solution in the presence of XXXII. The amount of OA solution consumed in the titration of the added quantity of Na₂S is determined separately. To samples XVI and XVII, up to 20 ml 1 N NaOH solution is added, heated to boiling, and an excess of 0.05 N OA solution is added; after 5 min, 30 ml 1 N NH₄NO₃ solution, 50 ml cold water and 2 - 4 ml Na₂S solution are added, and the Na₂S excess is titrated in the presence of XXXI. XVIII - XX are boiled for 5 - 10 min in alkaline

Card 3/4

Desulfurating titration of organic ...

S/081/63/000/003/008/036
B144/B186

solution with CA excess. XXI - XXX cannot be determined by the method described. I - X can be determined in the presence of XIII - XXX; therefore titration must be conducted at 25°C. XXV - XXX do not interfere with the determination of I - XVII. [Abstracter's note: Complete translation.]

Card 4/4

WRONSKI, Mieczyslaw

Mercurimetric determination of some sulfides.
Nauki matemat. przyrod. Lodz no.13:141-145 '62.

1. Katedra Technologii Chemicznej, Uniwersytet, Lodz.

WRONSKI, Mieczyslaw

Determination of equivalent weight of organic acids by titration
of benzylthiuronium salts with a HMB solution. Chem anal
8 no.1:113-115 '63.

1. Katedra Technologii Chemicznej, Uniwersytet, Lodz.

WRONSKI, Mieczyslaw

Thiomercurimetric determination of boron organic compounds.
Chem anal 8 no.2:299-300 '63.

1. Katedra Technologii Chemicznej, Uniwersytet, Lodz.

WRONSKI, Mieczyslaw

Mercurimetric determination of cystine together with
cysteine and sulfides. Chem anal 8 no.3:467-471 '63.

1. Katedra Technologii Chemicznej, Uniwersytet, Lodz.

WRONSKI, Mieczyslaw; BOGDANSKI, Janusz

Kinetics of cyanoethylation reaction of water, alcohols, amines,
and sulfhydryl compounds. Nauki matemat. przyrod. Lodz no.14:153-174
'63.

1. Katedra Technologii Chemicznej, Uniwersytet, Lodz.

WRONSKI, Mieczysław, doc. dr inż.

Thiomercurimetric titration. Wiad chem 17 no.1:1-27 Ja '63.

1. Kierownik Katedry Technologii Chemicznej, Uniwersytet,
Lodz.

WRONSKI, Mieczyslaw, doc. dr

Thiomercurometric determination of nitrites. Chem anal 9 no.1:
169-170 '64.

1. Katedra Technologii Chemicznej, Uniwersytet, Lodz.

L 31411-66
 ACC NR: AP6022964 SOURCE CODE: CZ/0008/65/000/009/1079/1085
 AUTHOR: Wronski, Mieczyslaw 25
 ORG: Institute of Chemical Technology, University, Lodz (Instytut Technologii B
 Chemicznej Uniwersytetu)
 TITLE: Analytical methods in the chemistry of sulfur compounds based on the
 application of mercury compounds
 SOURCE: Chemicke listy, no. 9, 1965, 1079-1085
 TOPIC TAGS: desulfurization, sulfide, mercaptan, cystine
 ABSTRACT: S compounds are usually determined by reactions based
 on neutralization, oxidation, or formation of complexes. For
 selective determination of S, its compounds with metals are used.
 The use of Cu and Ag is reviewed and the limitations of these
 metals discussed. Hg offers these advantages: The bond between
 Hg and S is very strong; mercurimetric titrations are suitable
 even in the presence of substances that would make other methods
 unusable. The equivalence point can be indicated electrometric-
 ally or by the use of indicators; a great number of organic com-
 pounds of mercury may be used as reagents. Selective determina-
 tions of sulfides and mercaptans, cystine and cysteamine, sel-
 ective desulfurization titration, and the use of selective mask-
 ing agents are discussed. Orig. art. has: 2 tables. [JPRS]
 SUB CODE: 07 / SUBM DATE: 11Jun64 / ORIG REF: 034 / OTH REF: 057
 Card 1/1 8T 0915 1048

WRONSKI, S.

CA

Effect of temperature on the intensity of x-rays reflected from various planes of zinc crystal. S. Wronski. *Acta Phys. Polon.* 7, 357-60 (1938) (in German).—The relative intensities of x-rays reflected from various crystallographic planes of metallic Zn were detd. by the Debye-Scherrer method at room temp. and at 567°K. From these measurements the following values were calcd. for the amplitudes of thermal oscillation of Zn atoms at room temp.: $\mu_1 = 0$ (parallel to the c-axis) = 0.127 Å.; $\mu_2 = 90$ (perpendicular to the c-axis) = 0.0734 Å. The characteristic temps. calcd. from these values are $\theta_1 = 200^\circ\text{K}$. and $\theta_2 = 367^\circ\text{K}$. resp. R. Jozefowicz

ASM-A-1 METALLURGICAL LITERATURE CLASSIFICATION

1ST AND 2ND ORDERS																										3RD AND 4TH ORDERS																									
1ST AND 2ND ORDERS																										3RD AND 4TH ORDERS																									
WRONSKI S.																										A-1																									
AC																																																			
Influence of temperature on the intensity of Röntgen rays reflected from different planes of the zinc crystal. S. WRONSKI (Acta Phys. Polon., 1939, 7, 357—366).—The relative intensities of reflection from different lattice planes of the Zn crystal have been measured at room temp. and 567° K. by the Debye-Scherrer method. At room temp. the amplitudes of heat oscillations of the Zn atom in the direction of and normal to the c-axis are calc. to be 0.127 and 0.0734 Å. Calc. vals. for the characteristic temperature of Zn are Θ_u , 200° K., Θ_l , 347° K.																																																			
O. D. S.																																																			
ASR-51A METALLURGICAL LITERATURE CLASSIFICATION																																																			
MATERIALS INDEX																										SUBJECT INDEX																									
1ST AND 2ND ORDERS																										3RD AND 4TH ORDERS																									
1ST AND 2ND ORDERS																										3RD AND 4TH ORDERS																									

POLAND/Physical Chemistry. Thermodynamics. Thermochemistry. B
Phase Transitions. Equilibria. Physico-Chemical
Analysis.

Abs Jour: Ref. Zhur. - Khimiya, No. 4, 1959, 10987

Authors : Ciborowski J., Wronski S.

Inst : Not given

Title : A Psychometric Chart for the System, Air - Ethyl
Acetate.

Orig Pub: Chem. stosow, 1958, 2, 147-152.

Abstract: On the basis of literature data, a psychometric
diagram was drawn for the system, air - ethyl
acetate. A disagreement between the psychometric
and adiabatic lines was discovered. A comparison
of some points, taken from this diagram, with a
few experimental results, previously obtained
(Mark I. G., Trans. Amer. Inst. Chem. Engrs., 1932,

Card 1/2

COUNTRY	: Poland	H-8
CATEGORY	:	
ABS. JOUR.	: RZKhim., No. 21 1959, No.	75402
AUTHOR	: Ciborowski, J. and Wronski, S.	
INST.	: Not given	
TITLE	: The Reduction of Sodium Sulfate with Hydrogen in Fluidized Beds	
ORIG. PUB.	: Przemysl Chem, 37, No 8, 520-522 (1958)	
ABSTRACT	<p>: The possibility of carrying out the reduction of Na_2SO_4 in fluidized beds at temperatures exceeding the melting point of the eutectic has been investigated. The reaction proceeds at low sulfate concentrations and at high hydrogen rates, assuring intensive mixing. The sulfate is reduced in 8 min when mixtures containing 5 and 7.5% sulfate are used and the grain size in the charge is 0.15-0.3 mm, in the presence of 1% iron (catalyst). The reduction is accompanied by an increase in the size of the grains as a result of agglomeration.</p> <p style="text-align: right;">From authors' summary</p>	

CARD: 1/1

CIBOROWSKI, Janusz; WRONSKI, Stanislaw

Investigation of sublimating condensation of naphthalene by mixing with fluidal charge. Chemia stosow 3 no.4:447-460 '59. (KEAI 9:6)

1. Zaklad Inzynierii Chemicznej Politechniki Warszawskiej i
Instytutu Chemii Ogolnej.
(Naphthalene)

WRONSKI, S.

5(2)

SOV/80-32-3-1/43

AUTHORS: Cyboronaki, F., Wronski, S.

TITLE: Reduction of Sodium Sulfate by Hydrogen in a Pseudo-Liquefied Layer (Vosstanovleniye sul'fata natriya vodorodom v pseudo-ozhizhennom sloye)

PERIODICAL: Zhurnal prikladnoy khimii, 1959, Vol XXXII, Nr 3, pp 473-477 (USSR)

ABSTRACT: Na_2S may be obtained by the reduction of Na_2SO_4 using hydrogen as reducing agent [Ref 1, 2]. An apparatus has been developed for this purpose (Figure 1). The experiments were carried out in two series: in homogeneous Na_2SO_4 and in a mixture of Na_2SO_4 and Na_2S . The reaction in the homogeneous substance proceeded in various stages at 62C, 640, 680 and 720 - 760°C. The final product contained 88 - 97% Na_2S . In the mixture hydrogen was introduced at the rate of 20 l/min. At low temperatures the sulfide yield was 88%, above 700°C 97%. An iron catalyst in the amount of 1% was used in the experiments. The consumption of hydrogen was only 5% under the most favorable conditions.

Card 1/2 There are 3 graphs, 1 diagram and 10 references, 3 of which

SOV/80-32-3-1/43

Reduction of Sodium Sulfate by Hydrogen in a Pseudo-Liquefied Layer

are Soviet, 3 German, 2 English, 1 Polish and 1 American.

ASSOCIATION: Kafedra protsessov i apparatov khimicheskoy tekhnologii Varshavskogo politekhnicheskogo instituta i instituta obshchey khimii (Chair of Processes and Apparatuses of Chemical Technology of the Warsaw Polytechnical Institute and the Institute of General Chemistry)

SUBMITTED: June 17, 1958

Card 2/2

CIBOROWSKI, Janusz; WRONSKI, Stanislaw

Testing of sublimating condensation of naphthalene by mixing with a fluidal charge. Chemia stosow 3 no.4:447-460 '59.

1. Zaklad Inzynierii Chemicznej, Politechnika, Warszawa i Instytut Chemii Ogolnej, Warszawa.

CIBOROWSKI, Janusz; WRONSKI, Stanislaw

The continuous method of sublimating condensation in fluidised bed.
Przem chem 40 no.8:433-436 Ag '61.

1. Katedra Inzynierii Chemicznej Politechniki Warszawskiej i Zaklad
Inzynierii Chemicznej Instytutu Chemi Organicznej.

CIBOROWSKI, Janusz; WRONSKI, Stanislaw

Sublimating condensation in a membrane cooled fluidized bed.
Chemia stosow 6 no.2:153-165 '62.

1. Katedra Inzynierii Chemicznej, Politechnika, i Zaklad Inzynierii
Chemicznej, Instytut Chemii Ogolnej, Warszawa.

CIBOROWSKI, Janusz; WRONSKI, Stanislaw

Mass and heat transfer from fluidized bed of sublimate material to the cooler wall. Chemia stosow 6 no. 4:529-540 '62.

1. Katedra Inzynierii Chemicznej, Politechnika, Warszawa, i Zaklad Inzynierii Chemicznej, Instytut Chemii Ogolnej, Warszawa.

CIBOROWSKI, Janusz; WRONSKI, Stanislaw

Studies on the efficiency of heat recovery in a cyclone exchanger working with a fluidized-solid furnace. Przem chem 42 no.1:38-41 Ja '63.

1. Katedra Inzynierii Chemicznej, Politechnika, Warszawa.

WRONSKI, W.

Characteristics of casein fibers, p. 232. (PRZEMYSŁ WŁOKIENNYCZY, Lodz, Vol. 7, no. 9/10, Sept./Oct. 1953.)

SO: Monthly List of East European Accessions, (EEAL), LC, Vol. 4, No. 4, Jan. 1955, Uncl.

WROŃSKI, WT.

Swelling of casein fibers. Wl. Wroński and H. Jabłoński, *Przemysł Włókienniczy* 9, 239-42(1955).—The swelling of various types of fibers in the presence of water was discussed in general. The centrifugal method of detg. the swelling rate (SR) is described. It was established that the SR value of casein fibers depends on the compds. and temp. of the hardening bath as well as on stretching conditions. The presence of Al sulfate reduces the SR. SR does not depend on the HCHO concn. within the range of 25-40 g./l. It depends, however, upon the time and temp. of fixing. The presence of NaCl or Na₂SO₄ in the HCHO coagulating bath improves the water resistance of fibers; the effect of NaCl is slightly stronger than that of Na₂SO₄. In proportion to the increase in the degree of stretching SR decreases. It was also proved that deamination with a NaNO₂ soln. reduces the SR of casein fibers. The optimum coagulating temp. in relation to the swelling value was 68-70°. A. Wielopolski

2

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001961730003-1



APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001961730003-1"

WRONSKI, W.

POLAND/Chemical Technology - Chemical Products and Their
Application, Part 4. - Artificial and Synthetic
Fibers.

H-31

Abs Jour : Ref Zhur - Khimiya, No 7, 1958, 23449

Author : W. Wronski

Inst :

Title : Quality Problem of Artificial Protein Fibers.

Orig Pub : Przem. chem., 1957, 13, No 4, 199-204

Abstract : Bibliography with 19 titles.

Card 1/1

WRONSKI, Wieslaw, mgr

Size of deposit and profitableness of mining it. Rudy i metale
9 no.11:614-617 N '64.

WRONSKI, Wieslaw, mgr

Economic aspects of investing and management of mining copper deposits. Rudy i metale 8 no.7:254-257 Je '63.

WRONSKI, Wieslaw, mgr.; JARCZYK, Kazimierz, mgr

On the difficulties of practical application of the economic
indicator of investment effectiveness in mining. Rudy i
metale 7 no.8:367-369 Ag '62.

CYPRYK, Jerzy; WRONSKI, Włodzimierz

Coagulation of polyacrylonitrile solutions. Tworzywa wielkocząst 6
no.11:363-367 N '61.

S/081/62/000/024/020/052
B117/B186

AUTHORS: Cytryk, Jerzy, Wroński, Włodzimierz

TITLE: Coagulation of polyacrylonitrile solutions

PERIODICAL: Referativnyy zhurnal. Khimiya, no. 24 (II), 1962, 833-834,
abstract 24P95 (Polimery, tworzywa, wielkocząsteczkowa, v. 6,
no. 11, 1961, 363-367 [Pol.; summaries in Eng. and Russ.])

TEXT: The coagulation of polyacrylonitrile from aqueous solution of dimethyl formamide was studied. The effects due to temperature and concentration of dimethyl formamide, and those due to concentration of polyacrylonitrile solution, on the transparency of films was determined. Photographs are given showing the microstructures of films obtained at concentrations of a dimethyl formamide solution between 30 and 70 % at 20°C and at 60 % at 15, 30, and 40°C. It was shown that a transparent gel without bubbles forms from the 40 - 60 % aqueous dimethyl formamide solution below 20°C and at a concentration of polymer (molecular weight 73 000) \geq 20 %.
[Abstracter's note: Complete translation.]

Card 1/1

Wronski, Z.

TOWPIK. J.; WRONSKI, Z.

Numeric and clinical characteristics of late symptomatic syphilis.
Polski tygod. lek. 7 no. 17:519-526 28 Apr 1952. (CLML 22:4)

1. Of the Clinical Department (Head--J. Towpik, M. D.) of the National Institute of Dermatology and Venereology (Director--J. Suchanek, M. D.)

FONIOK, F.; WRONSKI, Z.

High-power ferrite resonance insulators. Pt. 1. Przem. inst telekom
prace 14 no.46:23-27 '64.

L L2019-65

P/2507/64/014/046/0023/0037

ACCESSION NR: AT5007776

AUTHOR: Foniok, F.; Wronski, Z. (Vron'ski, Z.)

TITLE: High-power ferrite resonance isolators. Part I. Design methods

SOURCE: Warsaw. Przemyslowy Instytut Telekomunikacji. Prace, v. 14, no. 46, 1964, 23-37

TOPIC TAGS: isolator design, ferrite isolator, resonance isolator, high power isolator, ferrite polarization, waveguide, dielectric loss, saturation magnetization

ABSTRACT: The article gives a comprehensive review of methods used in the design of high power ferrite resonance isolators consisting of ferrite and dielectric plates mounted in a waveguide. The most important design specifications are discussed. The reverse attenuation is also discussed.

power, P_{imp} , and upper and lower bounds
Card 1/4

L L2019-65

ACCESSION NR: AT5007776

that magnetization and for its temperature variations are given. The methods for choosing
the dielectric material are also discussed. Techniques for calculating the physical dimen-
sions of the dielectric plates are discussed in detail. The thickness h of the plates is
determined by the required capacitance C and the dielectric constant ϵ_r of the material.

tables.

Card 2/4

L 42019-65

ACCESSION NR: AT5007776

ASSOCIATION: Przemyslowy Instytut Telekomunikacji, Warsaw (Telecommunications Research Institute)

SUBMITTED: 26Oct63

ENCL: 01

SUB CODE: EC

NO REF SOV: 002

OTHER: 010

WROTEK, Jerzy

Optimum approach in isoparametric processes. Chemia stosow A
9 no.1:41-49 '65.

1. Department of Chemical Technology of Warsaw University.
Submitted February 1, 1964.

MALAWSKI, Marek J.; WROTEK, Jerzy

A method of graphic analysis of the kinetics of a system of interdependent chemical reactions. I. Rocz chemii 34 no.5:1297-1306 '60. (BEAI 10:9)

1. Katedra Chemii Organicznej Uniwersytetu, Warszawa.

(Chemical reactions)

WROTEK, Jerzy, mgr inz.

New joints for electric overhead conductors. Przegl kolej
elektrotech 11 [i.e. 16] no.5:153-154, My '64.

WROTNOWSKA, Barbara

Hydrogeological picture of the Chmielnik region. Kwartalnik
geol 5 nb.4:975-976 '61.

1. Zaklad Hydrogeologii, Instytut Geologiczny, Warszawa.

WROTON, L.

Pneumatic devices in machine tools. Pt. 1. (To be Contd) p. 141

PRZEGLAD MECHANICZNY. (Stowarzyszenie Inzynierow i Technikow Mechanikow
Polskich) Warszawa, Poland
Vol. 18, no.5, Mar. 1959

Monthly list of East European Accessions (EEAI) LC, Vol.8, no.7, July 1959

Uncl.

WROTHNY, L.

Pneumatic devices in machine tools. Pt. 2. p. 177

PRZEGLAD MECHANICZNY. (Stowarzyszenie Inzynierow i Technikow Mechanikow Polskich)
Warszawa, Poland
Vol. 18, no. 6, Mar. 1959

Monthly List of East European Accession (EEAI) LC, Vol. 8, no. 7, July, 1959

Uncl.

WROTONY, L.

Hydropneumatic Drives and Their Application in Machine Tools, Pt. 2, p. 276

PRZEGLAD MECHANICZNY (Stowarzyszenie Inzynierow i Technikow Mechanikow
Polskich)
Warszawa, Poland
Vol. 18, no. 9, May 1959

Monthly List of East European Accessions Index (EEAI), LC, Vol. 8, No. 11,
November 1959.

Uncl.

WROTNY, Lucjan Tadeusz, mgr., inż.

Standardization of feeds and spindle speeds in machine tools.
Mechanik 34 no.1:6-10 '62.

1. Politechnika Warszawska.

WROTONY, Lucjan Tadeusz, mgr., inż.

Standardization of spindle and feed speeds in machine tools.
Mechanik 35 no.2:61-63 '62.

1. Politechnika Warszawska, Członek komitetu redakcyjnego
miesięcznika "Mechanik"

WROTHNY, L.T., mgr inz.

"Tools and tool machines for machining metals" by D.H. Bruins."
Reviewed by L.T. Wrotny. Mechanik 35 no.10:584-585 0 '62.

WROTONY, L.T., mgr inz.

Soviet proposal for the classification of machine tools in five classes of precision. Mechanik 35 no.12:681-682 D '62.

1. Komisja Obrabiarek i Narzedzi, Polski Komitet Normalizacyjny, Warszawa.

WROTONY, Lucjan Tadeusz, mgr inż.

The need for extended research on the precision of machine tools.
Mechanik 36 no.1:43-46, 47 '63.

WROTOCHWIL-SKRZYPKOWA, Maria

Case of helminthiasis with manifestations of meningitis and Loeffler's syndrome. *Pediat. polska* 31 no.3:325-330 Mar 56.

1. Kliniki Chorob Zakaznych Wiek Dzieciacego A.M. w Warszawie
Kierownik: prof. dr. med. J. Bogdanowicz, Warszawa, ul. Wolska
37. Kliniki Chorob Zakaznych Wiek Dzieciacego.

(ASCARIASIS, in infant and child,
with Loeffler's synd. & meningitis (Pol))

(MENINGITIS, in infant and child,
in ascariasis, with Loeffler's synd. (Pol))

(LOEFFLER'S SYNDROME, in infant and child,
in ascariasis, with meningitis (Pol))

MIERZEJEWSKI, T.; MIRECKI, L.; PENSON, J.; SWICA, S.; WAJDA, Z.;
WROZOLKOWA, T.

Pheochromocytoma. Diagnostic value of aorto-arteriography and
therapeutic problems. Kardiol. pol. 6 no.3:155-159 '63.

1. Z II Kliniki Chorob Wewnętrznych Kierownik: prof. dr J.
Penson z Kliniki Radiologii i Radioterapii Kierownik: prof.
dr W. Grabowski z III Kliniki Chirurgicznej Kierownik: prof. dr
Z. Kieturakis z Zakładu Anatomii Patologicznej AM w Gdansk
Kierownik: prof. dr W. Czarnocki.
(PHEOCHROMOCYTOMA) (ANGIOGRAPHY)

WROZOSEK, A.

WROZOSEK, A. An appraisal of the up-to-date achievements of Polish industrial geography. p. 487.

Vol. 28, no. 3, 1956
PRZEGLAD GEOGRAFICZNY
Poland

So: East European Accession, Vol. 6, No. 5, May 1957

JURCZYK, Witold, dr. med.; STENGERT, Krzysztof; GARSTKA, Jerzy; TOKARZ, Feliks; WOLOWICKA, Laura; WRUK, Marian, dr.

Measures for the restoration of suddenly arrested blood circulation. Pol. tyg. lek. 20 no.10:354-356 8 Mr '65

1. Z Zakładu Anestezjologii przy III Katedrze Chirurgii Akademii Medycznej w Poznaniu (Kierownik Zakładu: dr. med. Witold Jurczyk; Kierownik Katedry: prof. dr. med. Adam Piskorz); z Katedry I Kliniki Prtopedycznej Akademii Medycznej w Poznaniu (Kierownik: prof. dr. Wiktor Dega) ; z Katedry i Kliniki Neurochirurgii Akademii Medycznej w Poznaniu (Kierownik: doc. dr. med. Hieronim Powiertowski) i z Oddziału Torakochirurgicznego Sanatorium PKP w Chodzieży (Ordynator Oddziału: dr. Marian Wruk; Dyrektor Sanatorium: dr. Ryszard Raczynski).

WRUK, Marian; ZBIGNIEW, Jasinski

Avulsion of the primary bronchus as a complication of severe trauma of the thorax. Polski przegl. chir. 30 no.5:576-578 May 58.

(THORAX, wds. & inj.

causing bronchial avulsion (Pol))

(BRONCHI, wds. & inj.

avulsion (Pol))

Wrzecionko J.

Character of alternating potential between electron and photon. J. Wrzecionko. *Bull. acad. polon. sci., Classe II 4, 000-74 (1958)*.—Electron-photon scattering according to the Tamm-Dancoff method is discussed; the electron-photon potential is calcd. according to the Dyson approxn. The angle variables and spin states are sepd.; the potential is studied for each spin state. This results in a state analogous to the nucleon- π particle potential. From *C.Z.* 1958, 1374. Henry M. Koehler

2